(11) Publication number: 0 626 402 A3

(12)

## **EUROPEAN PATENT APPLICATION**

(21) Application number: 94303811.7

(51) Int. Cl.5: C08G 61/08

(22) Date of filing: 26.05.94

(30) Priority: 27.05.93 US 68236 27.05.93 US 68240 15.12.93 US 167668

(43) Date of publication of application: 30.11.94 Bulletin 94/48

(84) Designated Contracting States: BE DE ES FR GB IT NL

Date of deferred publication of search report: 27.09.95 Bulletin 95/39

(71) Applicant: AMOCO CORPORATION 200 East Randolph Drive Chicago Illinois 60601 (US)

(72) Inventor: Nubel, Philip O. 819 Tulip Lane Naperville, Illinois 60540 (US)

Inventor: Yokelson, Howard B. 1339 Brush Hill Circle Naperville, Illinois 60540 (US) Inventor: Moreland, Robert B. 29W225 Hartmand Drive Naperville, Illinois 60564 (US) Inventor: Bagheri, Vahid 6000 Oakwood Drive, 5H Lisle, Illinois 80532 (US) Inventor: Cohen, Steven A. 962 East Prarie Avenue Naperville, Illinois 60540 (US) Inventor: Bouslog, William G. 1517 West Jefferson Naperville, Illinois 60540 (US) Inventor: Behrends, Raymond T. 425 North Stewart Avenue Lombard, Illinois 60148 (US) Inventor: Nelson, James P. 1121 Kings Point Court Naperville, Illinois 60563 (US)

(74) Representative : Schlich, George William et al Mathys & Squire 100 Grays Inn Road London WC1X 8AL (GB)

- (54) Process for preparing linear monofunctional and telechelic difunctional polymers and compositions obtained thereby.
- This invention relates to a process for preparation of non-crosslinked linear monofunctional and telechelic difunctional unsaturated polymers wherein the functional groups are reactive terminal groups other than vinyl groups. The average functionality number of the monofunctional unsaturated polymers is at least 0.7, as determined by nuclear magnetic resonance spectroscopy (NMR). The average functionality number of the telechelic difunctional polymers is at least 1.6-1.7, as determined by NMR. Monofunctional olefins and difunctional olefins are reacted with cyclic olefins or unsaturated polymers to prepare difunctional polymers. The process is substantially free of side reactions comprising double bond migration and cyclization. The unsaturated functional polymers can be reacted in condensation, addition and transesterification reactions to prepare polyesgraft ters. copolymers, polyurethanes, polyureas, polyamides, thermoplastic and therresins, block copolymers, exchange resins, adhesives, films, fibers, foams and flocculants.



## EUROPEAN SEARCH REPORT

EP 94 30 3811

Category	Citation of document with indication of relevant passages	n, where appropriate,	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Inc.)
A	EP-A-0 173 473 (BRITISH 5 March 1986 * page 12, line 27 - li	•		C08G61/08
A	EUROPEAN POLYMER JOURNA vol. 25, no. 7/8, 1989 pages 805-811, H.T.HO ET ALL. METATH OF SUBSTITUTED NORBORNE OF THE POLYMERS OF SOME * page 806; table 2 *	OXFORD GB, ESIS POLYMERIZATION NES:MICROSTRUCTURE	<b>-</b> -	
<b>A</b>	MACROMOLECULES, vol. 20, 1987 EASTON Un pages 778-782, P.A.PATTON ET ALL. 'FAI THE RATE OF TUNGSTEN HEXACHLORIDE/TETRAMETHY! OLEFIN METHATHESIS POLY! RING STRAIN IS NOT IMPO! * page 1, paragraph 1 *	CTORS INFLUENCING LTIN-CATALYSED MERIZATION:MONOMER	j jo namene.	TECHNICAL FIELDS
	page 1, paragraph 1	<b></b>	•	SEARCHED (Int.CL5)
				C08F
į			÷	
				·
	•			
	•			
	The present search report has been draw	wn up for all claims		
	Place of search	Date of completies of the search		Exception
	THE HAGUE	25 July 1995	Sti	enon, P
X: part Y: part	CATEGORY OF CITED DOCUMENTS  culturly relevant if taken alone culturly relevant if combined with another ment of the same category	I : theory or principle E : earlier patent door after the filing da D : document cited in L : document cited for	underlying the ument, but publi te the application	invention ished on, or

O BORN (4M M 12 O